

Joint Trauma System



Amputation: Evaluation and Treatment

Part of the Joint Trauma System (JTS) Clinical Practice Guideline (CPG) Training Series



Purpose



The CPG strives to provide standardization of optimal care for the performance of wound management and life-saving amputations that will ensure preservation of maximum limb length, promote healing of viable tissues, and facilitate optimal rehabilitative function.

This presentation is based on the [JTS Amputation: Evaluation and Treatment CPG, 01 Jul 2016 \(ID:07\)](#). It is a high-level review. Please refer to the complete CPG for detailed instructions. Information contained in this presentation is only a guideline and not a substitute for clinical judgment.

Agenda



1. Summary
2. Background
3. Evaluation
4. Decision Criteria
5. Amputation Prep
6. Precautions
7. Amputation Expectations
8. Amputation
9. Post Operative Management
10. Performance Improvement (PI) Monitoring
11. References
12. Appendices: N/A
13. Contributors

Summary



- Amputation may be required as a damage-control procedure in a massively injured patient.
- Intact or ability to restore perfusion can delay decision to amputate.

Background

Amputation Terminology

- **Traumatic Amputation:** Immediate extremity amputation caused by the wounding mechanism.
- **Primary Amputation:** Performed by a surgical team after evaluation of the mangled extremity and deciding not to pursue limb salvage.
- **Secondary Amputation:** Occurs after an initial attempt at limb salvage.
 - ❑ Early: Within 90 days
 - ❑ Late: After 90 days



Primary amputation performed at Role 2

Evaluation

- Evaluation of extremity begins with thorough inspection of the wound and perfusion and control of active hemorrhage.
 - ❑ May require surgical wound extension to inspect all levels of tissue.
 - ❑ Doppler and diagnostic arteriography are adjuncts to evaluation of perfusion.
- Gross decontamination and meticulous sharp debridement of non-viable tissue for all war wounds (see *War Wounds CPG*).



Traumatically amputated right lower extremity undergoing debridement

Decision Criteria



- When deciding to amputate, timing hinges on vascularity of the injured extremity. Consider the following:
 - ❑ Determine if intact or potential to restore perfusion by vascular repair or shunt.
 - ❑ If perfusion can be restored, decision to amputate for nerve or bone loss can be deferred until later.
 - ❑ Amputation may be necessary as a damage control procedure in a massively injured patient due to the amount of time required to restore perfusion.

Decision Criteria

- Ipsilateral fractures should be stabilized and should not impact your decision to amputate.
- Scoring systems to predict amputation need are not widely accepted or validated in the combat trauma population.



Complicated lower extremity with potential requirement for primary amputation

Decision Criteria

- Primary and early secondary amputations are most commonly performed for:
 - Vascular injury
 - Nerve injury not amenable to repair or functional extremity
 - Extensive loss or contamination of soft tissue
- Late secondary amputations are generally performed due to patient preference or major complications.

Prolonged damage control procedures for other injuries and patient instability resulted in prolonged ischemia of lower extremity necessitating early secondary amputation.



Amputation Prep

- Thorough inspection of all levels of tissue of the wound are required.
 - Extent of the zone of injury dependent on mechanism, treatments, and contamination load.
 - Tissue damage is often beyond that which is apparent on initial visual inspection.
- Control any active hemorrhage, debride non-viable tissue, and thoroughly irrigate wounds.



Heavily contaminated lower extremity after initial traumatic amputation

Precautions



- Accept atypical skin and tissue flaps as long as the tissue is viable.
- Do not perform primary closure of traumatic amputations.
 - All war wounds should be left open and re-evaluated with serial irrigation and debridement.
- Avoid open circular or guillotine amputations.
 - They sacrifice viable soft tissue and relegate the casualty to more proximal revisions.
 - Have not been shown to be significantly faster than length-preserving procedures.

Amputation Expectations

- Current consensus regarding extremity amputation following battle-injury is to:
 - ❑ Preserve limb length and vascularity.
 - ❑ Facilitate adequate wound drainage.
 - ❑ Achieve eventual coverage and closure of the amputation wound.

Atypical length and tissue flaps after amputation of battle-injury to lower extremity. Wound is left open to facilitate wound drainage.



Amputation


- If amputation is required, appropriate vascular structures should be ligated proximal to bone resection, but distal enough to allow healing.
 - ❑ Separate vascular structures from nerves prior to ligation.
- Amputations should be performed at the most distal level which provides viable bone and soft tissue for later closure.
 - ❑ If near proximal joint, preservation of bone length without soft tissue coverage advised to provide later options for reconstruction.
 - ❑ Re-evaluate amputation site within first 24 hours.



Slightly atypical soft tissue flaps with NPWT to facilitate drainage after irrigation and debridement.

Post-Operative Management

- Place soft dry dressings around the amputation site and extremity.
 - ❑ Circumferential wraps with gauze rolls and ace wraps in figure of eight fashion.
 - ❑ Avoid excessive compression.
- Place in splint or bivalve cast to prevent joint contracture and provide soft tissue support.
 - ❑ Make sure there is simple access for wound inspections.
- If short skin flaps, skin traction to prevent soft tissue traction is an option.
- Avoid placement of pillows under knees to prevent contractures.



Skin Traction

Emergency War Surgery Handbook,
4th Edition

Post-Operative Management

- Negative pressure wound therapy using reticulated open cell foam can be useful after complete wound debridement and hemostasis achieved.
 - ❑ Can be left in place for 24 to 48 hours.
 - ❑ Care to avoid occlusion and leak of seal is essential.
 - ❑ May macerate healthy tissue, obliterate soft tissue planes, and has a potential role in heterotopic ossification.
 - ❑ Problems include: bulky for transport; occlusion of tubing or leak; maceration of healthy tissue; and obliteration of soft tissue planes.



Gradual closure of extremity amputation wound after NPWT.

Post-Operative Management



- Coordinate all dressing changes and repeat debridement with evacuation schedule and plan to perform them in operating room.
 - OR provides access to equipment for unexpected issues.
 - OR provides anesthetic for patient comfort.

PI Monitoring



■ Intent (Expected Outcomes)

- All amputation wounds are appropriately dressed but NOT primarily closed in theater

■ Performance/Adherence Measures

- All amputation wounds are dressed but not closed in theater

■ Data Source

- Patient Record
- Department of Defense Trauma Registry (DoDTR)

References



1. Emergency War Surgery Handbook, 3rd United States Revision 2004. Borden Institute. Walter Reed Army Medical Center, Washington, D.C. Chapter 22: Soft Tissue Injuries. 2004: 22.1-22.14
2. Pollak AN. Use of Negative Pressure Wound Therapy with Reticulated Open Cell Foam for Lower Extremity Trauma. J Orthop Trauma 2008:
3. Schiro GR, Sessa S, Piccioli A, Maccauro G. Primary amputation vs limb salvage in mangled extremity: a systematic review of the current scoring system. BMC Musculoskelet Disord. 2015; 16:372.
4. Peck MA, Clouse WD, Cox MW, Bowser AN, Eliason JL, Jenkins DH, Smith DL, Rasmussen TE. The complete management of extremity vascular injury in a local population: A wartime report from the 332nd Expeditionary Medical Group/Air Force Theater Hospital, Balad Air Base, Iraq. J Vascular Surgery 2007; 45:1197-1205
5. Leininger, BE, Rasmussen TE, Smith DL, Jenkins DH, Coppola C. Experience with wound VAC and delayed primary closure of contaminated soft tissue injuries in Iraq. J Trauma 2006; 61: 1207-11
6. Gordon WT, O'Brien FP, Strauss JE, Andersen RC, Potter BK. Outcomes Associated With the Internal Fixation of Long Bone Fractures Proximal to Ipsilateral Traumatic Amputations. J Bone Joint Surg Am. 2010 Oct 6;92(13):2312-8.
7. Fox CJ, Gillespie DL, O'Donnell SD, Rasmussen TE, Goff JM, Johnson CA, et al. Contemporary management of wartime vascular injury. J Vasc Surg 2005;41:638-44
8. Clouse WD, Rasmussen TE, Peck MA, Eliason JL, Cox MW, Bowser AN, et al. In theater management of vascular injury: two years of the Balad Vascular Registry. J Am Col Surg 2007;204:625-32
9. Powell ET. The Role of Negative Pressure Wound Therapy with Reticulated Open Cell Foam in the Treatment of War Wounds. J Orthop Trauma 2008; 22: S138-S141
10. Webb, LX, Pape HP. Current Thought Regarding the Mechanism of Action of Negative Pressure Wound Therapy with Reticulated Open Cell Foam. J Orthop Trauma 2008; 22: 135-13

Contributors



- LtCol Wade Gordon, MC, USAF
- CDR Luke Balsamo, MC, USN
- LtCol Max Talbot, RCMS, CF
- LCDR Charles Osier, MC, USN
- LTC Anthony Johnson, MC, USA

- John Shero, MHA
- LTC Benjamin Potter, MC, USA
- CAPT Zsolt Stockinger, MC, USN

Slides: Maj Andrew Hall, MC, USAF
Images from JTS collection unless otherwise cited.